

## On $\phi$ -2-absorbing primary subsemimodules over commutative semirings

*Issaraporn Thongsomnuk, Ronnason Chinram  
Pattarawan Singavananda and Patipat Chumket*

**Abstract.** In this paper, we introduce the concepts of  $\phi$ -2-absorbing primary subsemimodules over commutative semirings. Let  $R$  be a commutative semiring with identity and  $M$  be an  $R$ -semimodule. Let  $\phi : S(M) \rightarrow S(M) \cup \{\emptyset\}$  be a function, where  $S(M)$  is the set of subsemimodules of  $M$ . A proper subsemimodule  $N$  of  $M$  is said to be a  $\phi$ -2-absorbing primary subsemimodule of  $M$  if  $rsx \in N \setminus \phi(N)$  implies  $rx \in N$  or  $sx \in N$  or  $rs \in \sqrt{(N : M)}$ , where  $r, s \in R$  and  $x \in M$ . We prove some basic properties of these subsemimodules, give a characterization of  $\phi$ -2-absorbing primary subsemimodules, and investigate  $\phi$ -2-absorbing primary subsemimodules of quotient semimodules.

### 1. Introduction

In 2007, the concept of 2-absorbing ideals of rings was introduced by Badawi [3]. He defined a *2-absorbing ideal*  $I$  of a commutative ring  $R$  to be a proper ideal and if whenever  $a, b, c \in R$  with  $abc \in I$ , then  $ab \in I$  or  $ac \in I$  or  $bc \in I$ . Later in 2011 [7], Darani and Soheilnia introduced the concept of 2-absorbing submodules and studied their properties. A proper submodule  $N$  of an  $R$ -module  $M$  is said to be a *2-absorbing submodule* of  $M$  if  $a, b \in R$  and  $m \in M$  with  $abm \in N$ , then  $am \in N$  or  $bm \in N$  or  $ab \in (N : M)$ .

In 2012, Chaudhari introduced the concept of 2-absorbing ideals of a commutative semiring in [6]. He defined a *2-absorbing ideal*  $I$  of a commutative semiring  $R$  to be a proper ideal and if whenever  $a, b, c \in R$  with  $abc \in I$ , then  $ab \in I$  or  $ac \in I$  or  $bc \in I$ . In the same year, Thongsomnuk

2010 Mathematics Subject Classification: 13C05, 13C13, 16Y60

Keywords: Semimodule,  $\phi$ -2-absorbing primary subsemimodule, subtractive subsemimodule, Q-subsemimodule

introduced the concept of *2-absorbing subsemimodules over commutative semirings* as a proper subsemimodule  $N$  of an  $R$ -semimodule  $M$  such that if whenever  $a, b \in R$  and  $m \in M$  with  $abm \in N$ , then  $am \in N$  or  $bm \in N$  or  $ab \in (N : M)$ . The concept of 2-absorbing ideals of commutative semirings and 2-absorbing subsemimodules has been widely recognized by several mathematicians, see [8] and [11].

Atani and Kohan (2010) introduced and examined the concept of primary ideals in a commutative semiring, as well as primary subsemimodules in semimodules over a commutative semiring (see [5]). They defined a *primary ideal  $I$  of a commutative semiring  $R$*  as a proper ideal, such that whenever  $a, b \in R$  with  $ab \in I$ , then  $a \in I$  or  $b^k \in I$  for some  $k \in \mathbb{N}$ . Similarly, a *primary subsemimodule  $N$  of an  $R$ -semimodule  $M$*  is defined as a proper subsemimodule, such that whenever  $a \in R$  and  $m \in M$  with  $am \in N$ , then  $m \in N$  or  $a^k \in (N : M)$  for some  $k \in \mathbb{N}$ . In 2015, Dubey and Sarohe [9] defined the concept of 2-absorbing primary subsemimodules of a semimodule  $M$  over a commutative semiring  $R$  with  $1 \neq 0$  which is a generalization of primary subsemimodules of semimodules. A proper subsemimodule  $N$  of a semimodule  $M$  is said to be a *2-absorbing primary subsemimodule of  $M$*  if  $abm \in N$  implies  $ab \in \sqrt{(N : M)}$  or  $am \in N$  or  $bm \in N$  for some  $a, b \in R$  and  $m \in M$ .

Anderson and Batanieh (2008) generalized the concept of prime ideals, weakly prime ideals, almost prime ideals,  $n$ -almost prime ideals and  $\omega$ -prime ideals of rings to  $\phi$ -prime ideals of rings with  $\phi$ , see in [1]. They defined a  *$\phi$ -prime ideal  $I$  of a ring  $R$*  with  $\phi$  be a proper ideal and if for  $a, b \in R$ ,  $ab \in I \setminus \phi(I)$  implies  $a \in I$  or  $b \in I$ . Later in 2016, Petchkaew, Wasanawichit and Pianskool [13] introduced the concept of  $\phi$ - $n$ -absorbing ideals which are a generalization of  $n$ -absorbing ideals. A proper ideal  $I$  of  $R$  is called a  *$\phi$ - $n$ -absorbing ideal* if whenever  $x_1, x_2, \dots, x_{n+1} \in I \setminus \phi(I)$  for  $x_1, x_2, \dots, x_{n+1} \in R$ , then  $x_1 x_2 \dots x_{i-1} x_{i+1} \dots x_{n+1} \in I$  for some  $i \in \{1, 2, \dots, n+1\}$ . In 2017, Moradi and Ebrahimpour [12] introduced the concept of  $\phi$ -2-absorbing primary and  $\phi$ -2-absorbing primary submodules. Let  $\phi : S(M) \rightarrow S(M) \cup \{\emptyset\}$  be a function, where  $S(M)$  is the set of  $R$ -module  $M$ . They said that a proper submodule  $N$  of  $M$  is a  *$\phi$ -2-absorbing primary submodule* if  $rsx \in N \setminus \phi(N)$  implies  $rx \in N$ , or  $sx \in N$ , or  $rs \in \sqrt{(N : M)}$ , where  $r, s \in R$  and  $x \in M$ .

In this paper, we extend the concepts of  $\phi$ -2-absorbing primary submodules over commutative rings to the concepts of  $\phi$ -2-absorbing primary subsemimodules over commutative semirings. We explore fundamental prop-

erties of these subsemimodules, provide a characterization of  $\phi$ -2-absorbing primary subsemimodules, and investigate  $\phi$ -2-absorbing primary subsemimodules of quotient semimodules.

## 2. Preliminaries

**Definition 2.1.** [10] Let  $R$  be a semiring. A *left  $R$ -semimodule* (or a left semimodule over  $R$ ) is a commutative monoid  $(M, +)$  with additive identity  $0_M$  for which a function  $R \times M \rightarrow M$ , denoted by  $(r, m) \mapsto rm$  and called the scalar multiplication, satisfies the following conditions for all elements  $r$  and  $r'$  of  $R$  and all elements  $m$  and  $m'$  of  $M$ :

- (1)  $(rr')m = r(r'm)$ ,
- (2)  $r(m + m') = rm + rm'$ ,
- (3)  $(r + r')m = rm + r'm$ ,
- (4)  $1_R m = m$ , and
- (5)  $r0_M = 0_M = 0_R m$ .

Throughout this paper, we assume that  $R$  is a commutative semirings identity  $1 \neq 0$  and a left  $R$ -semimodule will be considered as a unitary semimodule.

**Definition 2.2.** [10] Let  $M$  be an  $R$ -semimodule and  $N$  a subset of  $M$ . We say  $N$  is a *subsemimodule of  $M$*  precisely when  $N$  is itself an  $R$ -semimodule with respect to the operations for  $M$ .

**Definition 2.3.** [5] Let  $M$  be an  $R$ -semimodule,  $N$  a subsemimodule of  $M$ , and  $m \in M$ . Then an *associated ideal* of  $N$  is denoted as

$$(N : M) = \{r \in R \mid rM \subseteq N\} \text{ and } (N : m) = \{r \in R \mid rm \in N\}.$$

**Definition 2.4.** [5] An ideal  $I$  of a semiring  $R$  is called a *subtractive ideal* if  $a, a + b \in I$  and  $b \in R$ , then  $b \in I$ .

A subsemimodule  $N$  of an  $R$ -semimodule  $M$  is called a *subtractive subsemimodule* if  $x, x + y \in N$  and  $y \in M$ , then  $y \in N$ .

**Proposition 2.5.** [5] Let  $M$  be an  $R$ -semimodule. If  $N$  is a subtractive subsemimodule of  $M$  and  $m \in M$ , then  $(N : M)$  and  $(N : m)$  are subtractive ideals of  $R$ .

**Lemma 2.6.** *Let  $(N : M)$  be a subtractive ideal of  $R$ . If  $a \in (N : M)$  and  $a + b \in \sqrt{(N : M)}$ , then  $b \in \sqrt{(N : M)}$ .*

*Proof.* Assume that  $a \in (N : M)$  and  $a + b \in \sqrt{(N : M)}$ . There exists  $k \in \mathbb{N}$  such that  $(a + b)^k \in (N : M)$ . Then  $\sum_{i=0}^k \binom{k}{i} a^{k-i} b^i \in (N : M)$ . Since  $\sum_{i=0}^{k-1} \binom{k}{i} a^{k-i} b^i \in (N : M)$  and  $(N : M)$  is a subtractive ideal, we obtain  $b^k \in (N : M)$ . Thus,  $b \in \sqrt{(N : M)}$ .  $\square$

**Definition 2.7.** [12] Let  $M$  be an  $R$ -semimodule. We define the functions  $\phi_\alpha : S(M) \rightarrow S(M) \cup \{\emptyset\}$  as follows:  $\phi_0(N) = 0$ ,  $\phi_\emptyset(N) = \emptyset$ ,  $\phi_{m+1}(N) = (N : M)^m N$  for every  $m \geq 0$  and  $\phi_\omega(N) = \bigcap_{m=0}^{\infty} (N : M)^m N$ , where  $N$  is a subsemimodule of  $M$  and  $S(M)$  is the set of subsemimodules of  $M$ .

**Definition 2.8.** [12] Let  $M$  be an  $R$ -semimodule,  $S(M)$  the set of subsemimodules of  $M$  and let  $f_1, f_2 : S(M) \rightarrow S(M) \cup \{\emptyset\}$  be two functions. Then  $f_1 \leq f_2$  if  $f_1(N) \subseteq f_2(N)$  for all  $N \in S(M)$ .

**Definition 2.9.** [2] A subsemimodule  $N$  of an  $R$ -semimodule  $M$  is called a *partitioning subsemimodule* (or  *$Q$ -subsemimodule*) if there exists a nonempty subset  $Q$  of  $M$  such that

1.  $RQ \subseteq Q$  where  $RQ = \{rq | r \in R \text{ and } q \in Q\}$ ,
2.  $M = \cup\{q + N | q \in Q\}$  where  $q + N = \{q + n | n \in N\}$ , and
3. if  $q_1, q_2 \in Q$ , then  $(q_1 + N) \cap (q_2 + N) \neq \emptyset$  if and only if  $q_1 = q_2$ .

Let  $M$  be an  $R$ -semimodule and  $N$  a  $Q$ -subsemimodule of  $M$ . Let  $M/N_{(Q)} = \{q + N | q \in Q\}$ . Then  $M/N_{(Q)}$  is a semimodule over  $R$  under the addition  $\oplus$  and the scalar multiplication  $\odot$  defined as follow: for any  $q_1, q_2, q \in Q$  and  $r \in R$ ,  $(q_1 + N) \oplus (q_2 + N) = q_3 + N$  and  $r \odot (q + N) = q_4 + N$  where  $q_3, q_4 \in Q$  are the unique elements such that  $q_1 + q_2 + N \subseteq q_3 + N$  and  $rq + N \subseteq q_4 + N$ . The  $R$ -semimodule  $M/N_{(Q)}$  is called the *quotient semimodule of  $M$  by  $N$* .

**Lemma 2.10.** [4] *Let  $M$  be an  $R$ -semimodule,  $N$  a  $Q$ -subsemimodule of  $M$  and  $P$  a subtractive subsemimodule of  $M$  with  $N \subseteq P$ . Then the followings hold:*

1.  $N$  is a  $Q \cap P$ -subsemimodule of  $P$ .
2.  $P/N_{(Q \cap P)} = \{q + N | q \in Q \cap P\}$  is a subsemimodule of  $M/N_{(Q)}$ .

**Remark 2.11.** *The zero element of  $P/N_{Q \cap P}$  is the same as the zero element of  $M/N_{(Q)}$  which is  $0_M + N$ .*

### 3. $\phi$ -2-absorbing primary subsemimodules

In this section, we investigate the  $\phi$ -2-absorbing primary subsemimodules over commutative semirings. Initially, we introduce a novel definition for  $\phi$ -2-absorbing primary subsemimodules. Subsequently, we explore various properties of  $\phi$ -2-absorbing primary subsemimodules.

**Definition 3.1.** Let  $M$  be an  $R$ -semimodule,  $\phi : S(M) \longrightarrow S(M) \cup \{\emptyset\}$  a function, where  $S(M)$  is the set of subsemimodules of  $M$ . We say a proper subsemimodule  $N$  of  $M$  is a  $\phi$ -2-absorbing primary subsemimodule if whenever  $rsx \in N \setminus \phi(N)$  implies  $rx \in N$ , or  $sx \in N$ , or  $rs \in \sqrt{(N : M)} = \{a \in R \mid a^n M \subseteq N \text{ for some } n \in \mathbb{N}\}$ , where  $r, s \in R$  and  $x \in M$ .

**Theorem 3.2.** Let  $M$  be an  $R$ -semimodule,  $N$  a  $\phi$ -2-absorbing primary subsemimodule of  $M$  and  $K$  be a subsemimodule of  $M$  such that  $\phi(N \cap K) = \phi(N)$ . Then  $N \cap K$  is a  $\phi$ -2-absorbing primary subsemimodule of  $K$ .

*Proof.* Clearly,  $N \cap K$  is a proper subsemimodule of  $K$ . Let  $rsx \in (N \cap K) \setminus \phi(N \cap K)$  where  $r, s \in R$  and  $x \in K$ . We have  $rsx \in N \setminus \phi(N \cap K)$ . Thus,  $rsx \in N \setminus \phi(N)$  because  $\phi(N \cap K) = \phi(N)$ . Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$ , we obtain  $rx \in N$ , or  $sx \in N$ , or  $rs \in \sqrt{(N : M)}$ . If  $rx \in N$  or  $sx \in N$ , then  $rx \in N \cap K$  or  $sx \in N \cap K$  because  $x \in K$  and  $K$  is an  $R$ -semimodule. If  $rs \in \sqrt{(N : M)}$ , then  $(rs)^n M \subseteq N$  for some positive integer  $n$ . In particular,  $(rs)^n K \subseteq (rs)^n M \subseteq N$  and we know that  $(rs)^n K \subseteq K$ . Then  $(rs)^n K \subseteq N \cap K$  for some positive integer  $n$ . Thus,  $rs \in \sqrt{(N \cap K : K)}$ . Hence  $N \cap K$  is a  $\phi$ -2-absorbing primary subsemimodule of  $K$ .  $\square$

Consider the following example. Let  $R = \mathbb{Z}_0^+$  and  $M = \mathbb{Z}_0^+$ , where throughout this paper,  $\mathbb{Z}_0^+$  denotes the set of non-negative integers (including zero). We define the function  $\phi : S(\mathbb{Z}_0^+) \rightarrow S(\mathbb{Z}_0^+) \cup \{\emptyset\}$  by  $\phi(A) = \{0\}$  where  $A \in S(\mathbb{Z}_0^+)$ . Clearly,  $8\mathbb{Z}_0^+$  is a  $\phi$ -2-absorbing primary subsemimodule of  $\mathbb{Z}_0^+$  and  $m\mathbb{Z}_0^+$  is a subsemimodule of  $\mathbb{Z}_0^+$  where  $m \in \mathbb{Z}_0^+$ . We see that  $\phi(8\mathbb{Z}_0^+ \cap m\mathbb{Z}_0^+) = \{0\} = \phi(8\mathbb{Z}_0^+)$ . Then  $8\mathbb{Z}_0^+ \cap m\mathbb{Z}_0^+ = [8, m]\mathbb{Z}_0^+$  is a  $\phi$ -2-absorbing primary subsemimodule of  $m\mathbb{Z}_0^+$ . This example demonstrates the concept outlined in Theorem 3.13.

**Theorem 3.3.** Let  $M$  be an  $R$ -semimodule,  $\phi : S(M) \longrightarrow S(M) \cup \{\phi\}$  a function, and let  $N$  be a proper subsemimodule of  $M$ . Then the following conditions are equivalent:

1.  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$ .
2. For every  $r \in R$  and  $x \in M$  with  $rx \notin N$ ,

$$(N : rx) \subseteq (\sqrt{(N : M)} : r) \cup (N : x) \cup (\phi(N) : rx).$$

*Proof.* First, let  $a \in (N : rx)$ . Then  $arx \in N$ . If  $arx \in \phi(N)$ , then  $a \in (\phi(N) : rx)$ . If  $arx \notin \phi(N)$ , then  $arx \in N \setminus \phi(N)$ . Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  and  $rx \notin N$ , we have  $ax \in N$  or  $a \in (\sqrt{(N : M)} : r)$ . Hence  $(N : rx) \subseteq (\sqrt{(N : M)} : r) \cup (N : x) \cup (\phi(N) : rx)$ .

Conversely, let  $r, s \in R$  and  $x \in M$  with  $rsx \in N \setminus \phi(N)$  and  $rx \notin N$ . Since  $rsx \in N$  and  $rsx \notin \phi(N)$ , we obtain  $s \in (N : rx)$  and  $s \notin (\phi(N) : rx)$ . From  $(N : rx) \subseteq (\sqrt{(N : M)} : r) \cup (N : x) \cup (\phi(N) : rx)$ . Thus,  $s \in (\sqrt{(N : M)} : r)$  or  $s \in (N : x)$ . Hence,  $sr \in \sqrt{(N : M)}$  or  $sx \in N$ . Therefore,  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$ .  $\square$

Moradi and Ebrahimpour [12] introduced the definition of  $\phi$ -triple-zero within the context of submodules. In this work, we will extend and adapt this definition to apply specifically to subsemimodules.

**Definition 3.4.** Let  $M$  be an  $R$ -semimodule, and  $\phi : S(M) \longrightarrow S(M) \cup \{\emptyset\}$  a function. Assume that  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$ ,  $r, s \in R$  and  $x \in M$ . We say  $(r, s, x)$  is a  $\phi$ -triple-zero of  $N$  if  $rsx \in \phi(N)$ ,  $rx, sx \notin N$  and  $rs \notin \sqrt{(N : M)}$ .

**Theorem 3.5.** Let  $M$  be an  $R$ -semimodule,  $\phi : S(M) \longrightarrow S(M) \cup \{\emptyset\}$  a function, and let  $N$  be a subtractive subsemimodule of  $M$  such that  $\phi(N) \subseteq N$ . Assume that  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  and  $(r, s, x)$  is a  $\phi$ -triple-zero of  $N$ . Then the following statements hold:

1.  $r(N : M)x \subseteq \phi(N)$  and  $s(N : M)x \subseteq \phi(N)$ .
2.  $(N : M)^2x \subseteq \phi(N)$ .
3.  $rsN \subseteq \phi(N)$ .
4.  $r(N : M)N \subseteq \phi(N)$  and  $s(N : M)N \subseteq \phi(N)$ .

*Proof.* (1). Suppose that there exists  $t \in (N : M)$  such that  $rtx \notin \phi(N)$ . Since  $(r, s, x)$  is a  $\phi$ -triple-zero of  $N$ , we have  $rsx \in \phi(N)$ . So,  $r(s + t)x = rsx + rtx \notin \phi(N)$ . Since  $\phi(N) \subseteq N$ , we obtain  $r(s + t)x \in N \setminus \phi(N)$ .

Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  and  $rx, sx \notin N$ , we have  $r(t+s) \in \sqrt{(N:M)}$ . By Lemma 2.6 and  $rt \in (N:M)$ , we have  $rs \in \sqrt{(N:M)}$ , which is a contradiction with  $\phi$ -triple-zero of  $N$ . Therefore,  $r(N:M)x \subseteq \phi(N)$ . Similarly,  $s(N:M)x \subseteq \phi(N)$ .

(2). Suppose that there exists  $t, k \in (N:M)$  such that  $tkx \notin \phi(N)$ . Since  $(r, s, x)$  is a  $\phi$ -triple-zero of  $N$ , we have  $rsx \in \phi(N)$ . By part (1), we have  $stx, rtx \in \phi(N)$ . Thus,  $(t+r)(k+s)x \notin \phi(N)$ . Then  $(t+r)(k+s)x \in N \setminus \phi(N)$ . Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  and  $rx, sx \notin N$ , we have  $(t+r)(k+s) \in \sqrt{(N:M)}$ . By Lemma 2.6, we obtain  $rs \in \sqrt{(N:M)}$ , which is a contradiction with  $\phi$ -triple-zero of  $N$ . Hence,  $(N:M)^2x \subseteq \phi(N)$ .

(3). Suppose that there exists  $y \in N$  such that  $rsy \notin \phi(N)$ . Since  $(r, s, x)$  is a  $\phi$ -triple-zero of  $N$ , we have  $rsx \in \phi(N)$ . So,  $rs(x+y) \notin \phi(N)$ . Then  $rs(x+y) \in N \setminus \phi(N)$  because  $\phi(N) \subseteq N$ . Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule,  $r(x+y) \in N$  or  $s(x+y) \in N$  or  $rs \in \sqrt{(N:M)}$ . Since  $N$  is a subtractive subsemimodule and  $y \in N$ , we obtain  $rx \in N$  or  $sx \in N$  or  $rs \in \sqrt{(N:M)}$ , which is a contradiction with  $\phi$ -triple-zero of  $N$ . Therefore,  $rsN \subseteq \phi(N)$ .

(4). Suppose that there exists  $t \in (N:M)$  and  $y \in N$  such that  $rtx \notin \phi(N)$ . Since  $(r, s, x)$  is a  $\phi$ -triple-zero of  $N$ , we obtain  $rsx \in \phi(N)$ . By parts (1) and (3), we have  $rtx, rsy \in \phi(N)$ . So,  $r(s+t)(x+y) \notin \phi(N)$ . Since  $\phi(N) \subseteq N$  and  $y \in N$ , we get  $r(s+t)(x+y) \in N \setminus \phi(N)$ . Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule,  $r(x+y) \in N$  or  $(s+t)(x+y) \in N$  or  $r(s+t) \in \sqrt{(N:M)}$ . Since  $N$  is a subtractive subsemimodule and Lemma 2.6, we have  $rx \in N$  or  $sx \in N$  or  $rs \in \sqrt{(N:M)}$ , which is a contradiction with  $\phi$ -triple-zero of  $N$ . Hence,  $r(N:M)N \subseteq \phi(N)$ . Similarly,  $s(N:M)N \subseteq \phi(N)$ .  $\square$

**Corollary 3.6.** *Let  $M$  be an  $R$ -semimodule,  $\phi : S(M) \longrightarrow S(M) \cup \{\emptyset\}$  a function, and let  $N$  be a subtractive subsemimodule of  $M$  such that  $\phi(N) \subseteq N$ . Assume that  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  and is not a 2-absorbing primary subsemimodule. Then  $(N:M)^2N \subseteq \phi(N)$ .*

*Proof.* Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  and is not a 2-absorbing primary subsemimodule, we have  $(r, s, x)$  is a  $\phi$ -triple-zero of  $N$ . Assume that  $t, k \in (N:M)$ ,  $y \in N$  and  $tky \notin \phi(N)$ . So,  $tky \in N \setminus \phi(N)$ . Consider  $(r+t)(s+k)(x+y) \notin \phi(N)$  because  $N$  is a  $\phi$ -triple zero and Theorem 3.5 and  $\phi(N) \subseteq N$  is subtractive subsemimodule. Then  $(r+t)(s+k)(x+y) \in N \setminus \phi(N)$ . Since  $N$  is a  $\phi$ -2-absorbing primary

subsemimodule, we have  $(r+t)(x+y) \in N$  or  $(s+k)(x+y) \in N$  or  $(r+t)(s+k) \in \sqrt{(N:M)}$ . Since  $N$  is a subtractive subsemimodule and Lemma 2.6, we have  $rx \in N$  or  $sx \in N$  or  $rs \in \sqrt{(N:M)}$ , which is a contradiction with  $\phi$ -triple-zero of  $N$ . Therefore,  $(N:M)^2N \subseteq \phi(N)$ .  $\square$

To illustrate Theorem 3.16(3), consider the following example. We define a function  $\phi : S(\mathbb{Z}_0^+) \rightarrow S(\mathbb{Z}_0^+) \cup \{\emptyset\}$  by  $\phi(A) = 2A$  where  $A \in S(\mathbb{Z}_0^+)$ . In this context,  $15\mathbb{Z}_0^+$  is demonstrably a  $\phi$ -2-absorbing primary subsemimodule and a subtractive subsemimodule of  $\mathbb{Z}_0^+$ . Interestingly,  $30\mathbb{Z}_0^+ = \phi(15\mathbb{Z}_0^+) \subseteq 15\mathbb{Z}_0^+$ . Furthermore, the triplet  $(3, 10, 2)$  qualifies as a  $\phi$ -triple-zero of  $15\mathbb{Z}_0^+$ . In this case,  $(3 \cdot 10) \cdot 15\mathbb{Z}_0^+ = 450\mathbb{Z}_0^+ \subseteq 30\mathbb{Z}_0^+$ , which aligns with the concept outlined in Theorem 3.16(3).

In 2017, the concept of weakly  $\phi$ -2-absorbing primary submodules was introduced by Moradi and Ebrahimpour [12]. In the current study, we will extend this idea and provide a definition for weakly  $\phi$ -2-absorbing primary subsemimodules.

**Definition 3.7.** Let  $M$  be an  $R$ -semimodule,  $\phi : S(M) \rightarrow S(M) \cup \{\emptyset\}$  be a function, where  $S(M)$  is the set of  $R$ -module  $M$ . They said that a proper submodule  $N$  of  $M$  is a *weakly  $\phi$ -2-absorbing primary submodule* if  $0 \neq rsx \in N \setminus \phi(N)$  implies  $rx \in N$ , or  $sx \in N$ , or  $rs \in \sqrt{(N:M)}$ , where  $r, s \in R$  and  $x \in M$ .

**Proposition 3.8.** Let  $M$  be an  $R$ -semimodule,  $\phi : S(M) \rightarrow S(M) \cup \{\emptyset\}$  a function, and let  $N$  be subtractive subsemimodule of  $M$  such that  $\phi(N) \subseteq N$  that is not 2-absorbing primary subsemimodule of  $M$ . If  $N$  is a weakly 2-absorbing primary subsemimodule of  $M$ , then  $(N:M)^2N = \{0\}$ .

*Proof.* Assume that  $N$  is a weakly 2-absorbing primary subsemimodule of  $M$  but  $N$  is not 2-absorbing primary subsemimodule of  $M$ . Then  $N$  is a  $\phi_0$ -2-absorbing primary subsemimodule of  $M$ . By Corollary 3.6, we obtain  $(N:M)^2N \subseteq \phi_0(N) = \{0\}$ . Clearly,  $\{0\} \subseteq (N:M)^2N$ . Thus,  $(N:M)^2N = \{0\}$ .  $\square$

Subsequently, we analyze the function  $\phi_n$ , as defined in Definition 2.7, for cases where  $n \leq 4$ . We also explore the function  $\phi_\omega$ , also defined in Definition 2.7, which establishes a connection with  $\phi$ -2-absorbing primary subsemimodules.

**Proposition 3.9.** Let  $M$  be an  $R$ -semimodule,  $\phi : S(M) \rightarrow S(M) \cup \{\emptyset\}$  a function, and let  $N$  be subtractive subsemimodule of  $M$  such that



$\phi(N) \subseteq N$  that is not 2-absorbing primary subsemimodule of  $M$ . If  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  for some  $\phi$  with  $\phi \leq \phi_4$ , then  $(N : M)^2N = (N : M)^3N$ .

*Proof.* Assume that  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  with  $\phi \leq \phi_4$  and  $N$  is not 2-absorbing primary subsemimodule. By Corollary 3.6, we obtain  $(N : M)^2N \subseteq \phi(N)$ . Since  $\phi \leq \phi_4$ , then  $\phi(N) \subseteq \phi_4(N) = (N : M)^3N$ . Now, we have  $(N : M)^2N \subseteq (N : M)^3N$ . Since  $N$  is an  $R$ -semimodule, we have  $(N : M)^3N = (N : M)(N : M)^2N \subseteq (N : M)^2N$ . Therefore,  $(N : M)^2N = (N : M)^3N$ .  $\square$

**Corollary 3.10.** *Let  $M$  be an  $R$ -semimodule,  $\phi : S(M) \longrightarrow S(M) \cup \{\emptyset\}$  a function, and let  $N$  be subtractive subsemimodule of  $M$  such that  $\phi(N) \subseteq N$ . If  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  with  $\phi \leq \phi_4$ , then  $N$  is a  $\phi_\omega$ -2-absorbing primary subsemimodule of  $M$ .*

*Proof.* Assume that  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  with  $\phi \leq \phi_4$ . It's clear that  $N$  is a  $\phi_\omega$ -2-absorbing primary subsemimodule of  $M$  if  $N$  is a 2-absorbing primary subsemimodule. Now, we consider in case that  $N$  is not 2-absorbing primary, then  $(N : M)^2N = (N : M)^3N$ , by Proposition 3.9. Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  with  $\phi \leq \phi_4$ , we have  $N$  is  $\phi_4$ -2-absorbing primary. So,  $\phi_\omega(N) = \bigcap_{m=0}^{\infty} (N : M)^mN = (N : M)^3N = \phi_4$ . Thus,  $N$  is a  $\phi_\omega$ -2-absorbing primary subsemimodule of  $M$ .  $\square$

**Lemma 3.11.** *Let  $N$  be a subtractive  $\phi$ -2-absorbing primary subsemimodule of an  $R$ -semimodule  $M$  and  $a, b \in R$ . Suppose that  $abK \subseteq N \setminus \phi(N)$  for some subsemimodule  $K$  of  $M$ . Then  $ab \in \sqrt{(N : M)}$  or  $aK \subseteq N$  or  $bK \subseteq N$ .*

*Proof.* Let  $abK \subseteq N \setminus \phi(N)$  for some subsemimodule  $K$  of  $M$ . Assume that  $ab \notin \sqrt{(N : M)}$ ,  $aK \not\subseteq N$  and  $bK \not\subseteq N$ . Then  $ak_1 \notin N$  and  $bk_2 \notin N$  for some  $k_1, k_2 \in K$ . Since  $abk_1 \in N \setminus \phi(N)$ ,  $ab \notin \sqrt{(N : M)}$ ,  $ak_1 \notin N$  and  $N$  is a  $\phi$ -2-absorbing primary subsemimodule, we have  $bk_1 \in N$ . Since  $abk_2 \in N \setminus \phi(N)$ ,  $ab \notin \sqrt{(N : M)}$ ,  $bk_2 \notin N$  and  $N$  is a  $\phi$ -2-absorbing primary subsemimodule, we obtain  $ak_2 \in N$ . We know that  $ab(k_1 + k_2) \in N \setminus \phi(N)$  and  $ab \notin \sqrt{(N : M)}$ . Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule, we have  $a(k_1 + k_2) \in N$  or  $b(k_1 + k_2) \in N$ . If  $a(k_1 + k_2) \in N$ , then  $ak_1 \in N$  (as  $N$  is a subtractive), which is a contradiction. If  $b(k_1 + k_2) \in N$ , then  $bk_2 \in N$  (as  $N$  is a subtractive), which is a contradiction. Hence,  $ab \in \sqrt{(N : M)}$  or  $aK \subseteq N$  or  $bK \subseteq N$ .  $\square$

**Theorem 3.12.** *Let  $K$  be a subtractive subsemimodule of  $M$  and  $\sqrt{(K : M)}$  be a subtractive ideal of  $R$ . If  $K$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$ , then whenever  $IJN \subseteq K \setminus \phi(K)$  for some ideals  $I, J$  of  $R$  and a subsemimodule  $N$  of  $M$ , then  $IJ \subseteq \sqrt{(K : M)}$  or  $IN \subseteq K$  or  $JN \subseteq K$ .*

*Proof.* Let  $K$  be a  $\phi$ -2-absorbing primary subsemimodule of  $M$ . Assume that  $IJN \subseteq K \setminus \phi(K)$  for some ideals  $I, J$  of  $R$  and a subsemimodule  $N$  of  $M$ . Suppose that  $IJ \not\subseteq \sqrt{(K : M)}$ ,  $IN \not\subseteq K$  and  $JN \not\subseteq K$ . Then  $a_1N \not\subseteq K$  and  $b_1N \not\subseteq K$  for some  $a_1 \in I$  and  $b_1 \in J$ . Since  $a_1b_1N \subseteq K \setminus \phi(K)$ ,  $a_1N \not\subseteq K$ ,  $b_1N \not\subseteq K$  and Lemma 3.11, we have  $a_1b_1 \in \sqrt{(K : M)}$ . Since  $IJ \not\subseteq \sqrt{(K : M)}$ , we have  $a_2b_2 \notin \sqrt{(K : M)}$  for some  $a_2 \in I$  and  $b_2 \in J$ . Since  $a_2b_2N \subseteq K \setminus \phi(K)$  and  $a_2b_2 \notin \sqrt{(K : M)}$ , we have  $a_2N \subseteq K$  or  $b_2N \subseteq K$  by Lemma 3.11. Here three cases arise.

**Case I:** When  $a_2N \subseteq K$  but  $b_2N \not\subseteq K$ . Since  $a_1b_2N \subseteq K \setminus \phi(K)$ ,  $b_2N \not\subseteq K$  and  $a_1N \not\subseteq K$ , then by Lemma 3.11,  $a_1b_2 \in \sqrt{(K : M)}$ . We know that  $a_2N \subseteq K$  but  $a_1N \not\subseteq K$ , so  $(a_1 + a_2)N \not\subseteq K$  (as  $K$  is subtractive). Since  $(a_1 + a_2)b_2N \subseteq K \setminus \phi(K)$ ,  $b_2N \not\subseteq K$  and  $(a_1 + a_2)N \not\subseteq K$ , we have  $(a_1 + a_2)b_2 \in \sqrt{(K : M)}$  by Lemma 3.11. Since  $a_1b_2 \in \sqrt{(K : M)}$  and  $\sqrt{(K : M)}$  is subtractive, we have  $a_2b_2 \in \sqrt{(K : M)}$ , which is a contradiction.

**Case II:** When  $b_2N \subseteq K$  but  $a_2N \not\subseteq K$ . We can conclude similarly to Case I.

**Case III:** When  $a_2N \subseteq K$  and  $b_2N \subseteq K$ . Since  $b_2N \subseteq K$  and  $b_1N \not\subseteq K$ , we have  $(b_1 + b_2)N \not\subseteq K$ . Since  $a_1(b_1 + b_2)N \subseteq K \setminus \phi(K)$ ,  $(b_1 + b_2)N \not\subseteq K$  and  $a_1N \not\subseteq K$ , we get that  $a_1(b_1 + b_2) \in \sqrt{(K : M)}$  by Lemma 3.11. Since  $a_1b_1 \in \sqrt{(K : M)}$  and  $\sqrt{(K : M)}$  is subtractive, we conclude that  $a_1b_2 \in \sqrt{(K : M)}$ . Since  $a_2N \subseteq K$ ,  $a_1N \not\subseteq K$  and  $K$  is subtractive implies  $(a_1 + a_2)N \not\subseteq K$ . Since  $(a_1 + a_2)b_1N \subseteq K \setminus \phi(K)$ ,  $(a_1 + a_2)N \not\subseteq K$  and  $b_1N \not\subseteq K$ , we have  $(a_1 + a_2)b_1 \in \sqrt{(K : M)}$  by Lemma 3.11. Since  $a_1b_1 \in \sqrt{(K : M)}$ ,  $(a_1 + a_2)b_1 \in \sqrt{(K : M)}$  and  $\sqrt{(K : M)}$  is subtractive, we have  $a_2b_1 \in \sqrt{(K : M)}$ . Since  $(a_1 + a_2)(b_1 + b_2)N \subseteq K \setminus \phi(K)$ ,  $(a_1 + a_2)N \not\subseteq K$  and  $(b_1 + b_2)N \not\subseteq K$ , by Lemma 3.11,  $(a_1 + a_2)(b_1 + b_2) \in \sqrt{(K : M)}$ . Since  $a_2b_1, a_1b_2, a_1b_1 \in \sqrt{(K : M)}$  and  $\sqrt{(K : M)}$  is subtractive, then  $a_2b_2 \in \sqrt{(K : M)}$ , which is a contradiction.

Hence,  $IJ \subseteq \sqrt{(K : M)}$  or  $IN \subseteq K$  or  $JN \subseteq K$ .  $\square$

**Theorem 3.13.** *Let  $M$  an  $R$ -semimodule, and let  $\phi : S(M) \longrightarrow S(M) \cup \{\emptyset\}$  be a function. Assume that  $N$  is a subsemimodule of  $M$  such that  $\phi(N)$  is a*

2-absorbing primary subsemimodule of  $M$  and  $\phi(N) \subseteq N$ . Then  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  if and only if  $N$  is a 2-absorbing primary subsemimodule of  $M$ .

*Proof.* First, assume that  $N$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  and  $\phi(N)$  is a 2-absorbing primary subsemimodule of  $M$ . Let  $r, s \in R$  and  $x \in M$  with  $rsx \in N$ . Suppose that neither  $rx$  nor  $sx$  is in  $N$ . Here two cases arise.

**Case I:**  $rsx \in \phi(N)$ . Then  $rs \in \sqrt{(\phi(N) : M)} \subseteq \sqrt{(N : M)}$  because  $\phi(N)$  is a  $\phi$ -2-absorbing primary subsemimodule,  $\phi(N) \subseteq N$  and  $rx, sx \notin N$ .

**Case II:**  $rsx \notin \phi(N)$ . Since  $N$  is a  $\phi$ -2-absorbing primary subsemimodule and  $rx, sx \notin N$ , we obtain  $rs \in \sqrt{(N : M)}$ .

Conversely, it's clearly.  $\square$

Let  $M$  be an  $R$ -semimodule,  $N$  be a  $Q$ -subsemimodule of  $M$ . For a function  $\phi : S(M) \rightarrow S(M) \cup \{\emptyset\}$  we define the function  $\phi_N : S(M/N_{(Q)}) \rightarrow S(M/N_{(Q)}) \cup \{\emptyset\}$  by  $\phi_N(K/N) = \phi(K)/N_{(\phi(K) \cap Q)}$  if  $\phi(K) \neq \emptyset$ , and  $\phi_N(K/N) = \emptyset$  if  $\phi(K) = \emptyset$ , for every subsemimodule  $K$  of  $M$  with  $N \subseteq K$ .

**Theorem 3.14.** *Let  $M$  be an  $R$ -semimodule,  $N$  a  $Q$ -subsemimodule of  $M$  and  $P, \phi(P)$  are subtractive subsemimodules of  $M$  with  $N \subseteq P$ . Then  $P$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$  if and only if  $P/N_{(Q \cap P)}$  is a  $\phi_N$ -2-absorbing primary subsemimodule of  $M/N_{(Q)}$ .*

*Proof.* First, assume that  $P$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$ . Then we have  $P/N_{(Q \cap P)}$  is a subsemimodule of  $M/N_{(Q)}$ . Now let  $r, s \in R$  and  $q_1 + N \in M/N_{(Q)}$  where  $q_1 \in Q$  be such that  $rs \odot (q_1 + N) \in P/N_{(Q \cap P)} \setminus \phi_N(P/N_{(Q \cap P)})$ . Then there exists unique  $q_2 \in Q \cap P$  such that  $rs \odot (q_1 + N) = q_2 + N$  where  $rsq_1 + N \subseteq q_2 + N$ . Since  $q_2 \in P$  and  $N \subseteq P$ , we have  $rsq_1 + N \subseteq P$ . Since  $N \subseteq P$  and  $P$  is a subtractive subsemimodule,  $rsq_1 \in P$ . Since  $rsq_1 + N \subseteq q_2 + N \notin \phi_N(P/N_{(Q \cap P)})$ , we obtain  $rsq_1 + N \subseteq q_2 + N \notin \phi(P)/N_{(Q \cap \phi(P))}$ . Thus, we have  $rsq_1 = q_2 + x$  for some  $x \in N \subseteq \phi(P)$ . Since  $q_2 \notin Q \cap \phi(P)$ , we get  $q_2 \notin \phi(P)$ . Then  $rsq_1 = q_2 + x \notin \phi(P)$  because  $\phi(P)$  is subtractive. Now, we have  $rsq_1 \in P \setminus \phi(P)$ . Since  $P$  is a  $\phi$ -2-absorbing subsemimodule of  $M$ , it can be concluded that  $rq_1 \in P$  or  $sq_1 \in P$  or  $rs \in \sqrt{(P : M)}$ . We claim that  $r \odot (q_1 + N) \in P/N_{(Q \cap P)}$  or  $s \odot (q_1 + N) \in P/N_{(Q \cap P)}$  or  $rs \in \sqrt{(P/N_{(Q \cap P)} : M/N_{(Q)})}$ .

**Case I:**  $rq_1 \in P$ . Since  $q_1 \in Q$ , we have  $rq_1 \in Q$ . Then  $rq_1 \in Q \cap P$ . So,  $rq_1 + N \in P/N_{(Q \cap P)}$ . Moreover,  $r \odot (q_1 + N) = q_3 + N$  where  $q_3 \in Q$  is unique such that  $rq_1 + N \subseteq q_3 + N$ . Then  $rq_1 = q_3 + x_1$  for some  $x_1 \in N \subseteq P$ . Since  $P$  is subtractive, we have  $q_3 \in P$ . Thus,  $r \odot (q_1 + N) = q_3 + N \in P/N_{(Q \cap P)}$ .

**Case II:**  $sq_1 \in P$ . We can conclude similarly to Case I that  $s \odot (q_1 + N) \in P/N_{(Q \cap P)}$ .

**Case III:**  $rs \in \sqrt{(P : M)}$ . Then there exists  $k \in \mathbb{N}$  such that  $(rs)^k \in (P : M)$ . So,  $(rs)^k M \subseteq P$ . Let  $q + N \in M/N_{(Q)}$  where  $q \in Q$ . Consider  $(rs)^k \odot (q + N) = q_4 + N$  where  $q_4 \in Q$  is unique such that  $(rs)^k + N \subseteq q_4 + N$ . So,  $(rs)^k q = q_4 + x_2$  for some  $x_2 \in N \subseteq P$ . Since  $(rs)^k \in (P : M)$ , we have  $(rs)^k q \in P$ . Hence,  $q_4 \in P$  because  $P$  is subtractive. Then  $q_4 \in Q \cap P$ . Thus,  $(rs)^k \odot (q + N) = q_4 + N \in P/N_{(Q \cap P)}$ . Hence,  $rs \in \sqrt{(P/N_{(Q \cap P)} : M/N_{(Q)})}$ .

Therefore,  $P/N_{(Q \cap P)}$  is a  $\phi_N$ -2-absorbing primary subsemimodule of  $M/N_{(Q)}$ .

Conversely, assume that  $P/N_{(Q \cap P)}$  is a  $\phi_N$ -2-absorbing primary subsemimodule of  $M$ . Let  $r, s \in R$  and  $x \in M$  such that  $rsx \in P \setminus \phi(P)$ . Since  $N$  is a  $Q$ -subsemimodule of  $M$  and  $x \in M$ , we have  $x \in q_1 + N$  where  $q_1 \in Q$ . So,  $rsx \in rs \odot (q_1 + N)$ . Let  $rs \odot (q_1 + N) = q_2 + N$  where  $q_2$  is the unique element of  $Q$  such that  $rsq_1 + N \subseteq q_2 + N$ . Then  $rsx \in q_2 + N$ . So there is  $y \in N$  such that  $q_2 + y = rsx \in P$ . Since  $y \in N \subseteq P$  and  $P$  is subtractive, we obtain  $q_2 \in P$ . Then  $q_2 \in Q \cap P$ . Thus,  $rs \odot (q_1 + N) = q_2 + N \in P/N_{(Q \cap P)}$ . Consider  $rsx \notin \phi(P)$  and  $y \in N \subseteq \phi(P)$ . Since  $rsx = q_2 + y$  and  $\phi(P)$  is subsemimodule, we have  $q_2 \notin \phi(P)$  so that  $q_2 + N \notin \phi(P)/N_{(Q \cap \phi(P))} = \phi_N(P/N)$ . Now, we have  $rs \odot (q_1 + N) = q_2 + N \notin P/N_{(Q \cap P)} \setminus \phi_N(P/N)$ . Since  $P/N_{(Q \cap P)}$  is a  $\phi_N$ -2-absorbing primary subsemimodule of  $M/N_{(Q)}$ , we get  $r \odot (q_1 + N) \in P/N_{(Q \cap P)}$  or  $s \odot (q_1 + N) \in P/N_{(Q \cap P)}$  or  $rs \in \sqrt{(P/N_{(Q \cap P)} : M/N_{(Q)})}$ . Here three cases arise.

**Case I:**  $r \odot (q_1 + N) \in P/N_{(Q \cap P)}$ . Then  $r \odot (q_1 + N) = q_2 + N$  where  $q_2$  is the unique element of  $Q \cap P$  such that  $rq_1 + N \subseteq q_2 + N$ . Thus,  $rq_1 + N \subseteq q_2 + N \subseteq P$  because  $N \subseteq P$  and  $q_2 \in Q \cap P$ . So,  $x \in q_1 + N$  that  $rx \in r(q_1 + N) \subseteq rq_1 + N \subseteq q_2 + N \subseteq P$ . Thus,  $rx \in P$ .

**Case II:**  $s \odot (q_1 + N) \in P/N_{(Q \cap P)}$ . We can conclude similarly to Case I that  $sx \in P$ .

**Case III:**  $rs \in \sqrt{(P/N_{(Q \cap P)} : M/N_{(Q)})}$ . Then  $(rs)^k \odot M/N_{(Q)} \subseteq P/N_{(Q \cap P)}$  for some  $k \in \mathbb{N}$ . Let  $m \in M$ . So, there is unique  $q_3 \in Q$  such that  $m \in q_3 + N$  and  $(rs)^k m \in (rs)^k(q_3 + N) \subseteq (rs)^k \odot (q_3 + N) = q_4 + N$  where  $q_4$  is the unique element of  $Q$  such that  $(rs)^k q_3 + N \subseteq q_4 + N$ . Now,  $q_4 + N = (rs)^k \odot (q_3 + N) \in P/N_{(Q \cap P)}$ . Then  $(rs)^k m \in q_4 + N \subseteq P$ . So,  $(rs)^k M \subseteq P$ . Thus,  $(rs)^k M \subseteq P$ . Therefore,  $rs \in \sqrt{(P : M)}$ .

Hence,  $P$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$ .  $\square$

**Corollary 3.15.** *Let  $M$  be an  $R$ -semimodule,  $N$  a  $Q$ -subsemimodule of  $M$ , and let  $P$  and  $\phi(P)$  be subtractive subsemimodules of  $M$  with  $N \subseteq P$ . If  $\phi(P) = N$  and  $P$  is a  $\phi$ -2-absorbing primary subsemimodule of  $M$ , then  $P/N_{(Q \cap P)}$  is a weakly 2-absorbing primary subsemimodule of  $M/N_{(Q)}$ .*

*Proof.* Since  $\phi(P) = N$ , we have  $\phi_N(P/N) = \phi(P)/N = \{0\}$ . By Theorem 3.14, we conclude that  $P/N_{(Q \cap P)}$  is a weakly 2-absorbing primary subsemimodule of  $M/N_{(Q)}$ .  $\square$

## References

- [1] **D.D. Anderson and M. Batanieh**, *Generalizations of prime ideals*, Comm. Algebra, **36** (2008), 686 – 696.
- [2] **R.E. Atani, and S.E. Atani**, *On subsemimodules of semimodules*, Bul. Acad. Stiinte Repub. Mold. Mat., **63** (2010), no. 2, 20 – 30.
- [3] **A. Badawi**, *On 2-absorbing ideals of commutative rings*, Bull. Aust. Math. Soc., **75** (2007), no. 3, 417 – 429.
- [4] **J.N. Chaudhari and B.R. Bonde**, *On partitioning and subtractive subsemimodules of semimodules over semirings*, Kyungpook Math. J., **50** (2010), 329 – 336.
- [5] **J.N. Chaudhari and B.R. Bonde**, *Weakly prime subsemimodules of semimodules over semirings*, Int. J. Algebra, **5** (2011), no. 4, 167 – 174.
- [6] **J.N. Chaudhari**, *2-absorbing ideals in semirings*, Int. J. Algebra, **6** (2012), no. 6, 265 – 270.
- [7] **A.Y. Darani, and F. Soheilnia**, *2-absorbing and weakly 2-absorbing submodules*, Thai. J. Math., **9** (2011), no. 3, 577 – 584.
- [8] **M.K. Dubey and P. Sarohe**, *On 2-absorbing semimodules*, Quasigroups Related Syst., **21** (2013), 175 – 184.

- [9] **M. K. Dubey and P. Sarohe**, *On 2-absorbing primary subsemimodules over commutative semirings*, Bul. Acad. Stiinte Repub. Mold., Mat., **78** (2015), no. 2, 27 – 35.
- [10] **J.S. Golan**, *Semirings and their Applications*, Kluwer Academic Publishers, Dordrecht, (1999).
- [11] **P. Kumar, M.K. Dubey and P. Sarohe**, *On 2-absorbing ideals in commutative semiring*, Quasigroups Related Syst. **24** (2016), 67 – 74.
- [12] **R. Moradi and M. Ebrahimpour**, *On  $\phi$ -2-absorbing primary submodule*, Acta Math. Vietnam, **42** (2017), 27 – 35.
- [13] **P. Petchkaew, A. Wasanawichit, and S. Pianskool**, *Generalizations of  $n$ -absorbing ideals of commutative semirings*, Thai. J. Math., **14** (2016), no. 2, 477 – 489.

Received August 13, 2023

I. Thongsomnuk

Division of Mathematics, Faculty of Science and Technology, Phetchaburi Rajabhat University, Na Wung, Muang, Phetchaburi 76000, Thailand  
E-mail: issaraporn.tho@mail.pbru.ac.th

R. Chinram

Division of Computational Science, Faculty of Science, Prince of Songkla University, Hat Yai, Songkhla, 90110 Thailand  
E-mail: ronnason.c@psu.ac.th

P. Singavananda

Program in Mathematics, Faculty of Science and Technology, Songkhla Rajabhat University, Khoa-Roob-Chang, Muang, Songkhla 90000, Thailand  
pattarawan.pe@skru.ac.th

P. Chumket

Division of Mathematics, Faculty of Science Technology and Agriculture, Yala Rajabhat University, Tambol Sateng, Mueang, Yala 95000, Thailand  
E-mail: patipat.c@yru.ac.th